

# Stone duality for finitely valued algebras with a near-unanimity term

Marco Abbadini. University of Birmingham, UK.

Joint work with Adam Přenosil.

MOSAIC Workshop 2023, Vienna, Austria, 29 September 2023.

Goal: represent algebraic structures as algebras of continuous functions satisfying some local constraints.

Algebraic structure  $\longleftrightarrow$  Space + local constraints

1. Stone duality (1936): Boolean algebra  $\cong$  algebra of continuous functions from a Stone space  $X$  to  $\{0, 1\}$  with pointwise operations.  
Space: a Stone space.  
Local constraints: none.
2. Priestley duality (1969): bounded distributive lattice  $\cong$  algebra of order-preserving continuous functions from a Priestley space  $X$  to  $\{0 < 1\}$ . (Priestley space = Stone space with a partial order satisfying compatibility properties.)  
Space: a Stone space.  
Local constraints: the order.

Algebraic structure  $\longleftrightarrow$  Space + local constraints

We investigate the case where the **constraints** can be made *very* local: on subsets of cardinality  $\leq 2$ .  $\rightsquigarrow$  Easier to describe them.

$\mathbb{ISP}(\mathbf{L}) :=$  class of algebras **isomorphic** to a **subalgebra** of a **power** of  $\mathbf{L}$ .

Davey & Werner (1983): duality for classes of the form  $\mathbb{ISP}(\mathbf{L})$  where  $\mathbf{L}$  is a finite algebra with a near unanimity term. (E.g.:  $\mathbf{L}$  has a lattice reduct.)

Example: for  $\mathbf{L} = \{0, 1\}$  we get Stone/Priestley dualities.

Our main result: we give an analagous duality, where the generating algebra  $\mathbf{L}$  is allowed to be infinite.

Our motivation: representation of *positive MV-algebras*, which are algebras related to Łukasiewicz many-valued logic:  $\{0, 1\}$  is replaced by  $[0, 1]$ .

## Examples of generating algebra $\mathbf{L}$ :

- ▶  $\mathbf{L} = \{0, 1\}$ , with the signature of Boolean algebras.  $\rightsquigarrow$  Stone duality (1936).
- ▶  $\mathbf{L} = \{0, 1\}$ , with the signature of bounded distributive lattices.  $\rightsquigarrow$  Priestley duality (1969).
- ▶  $\mathbf{L} = [0, 1]$ , with the signature of MV-algebras.  $\rightsquigarrow$  Duality for weakly locally finite MV-algebras (Cignoli, Marra, 2012).
- ▶  $\mathbf{L} = [0, 1]$ , with the signature of positive MV-algebras, i.e.  $\{\oplus, \odot, \vee, \wedge, 0, 1\}$ .
- ▶  $\mathbf{L} = \mathbb{R}$ , with the signature of  $\ell$ -groups with a designated constant 1.
- ▶  $\mathbf{L} = \mathbb{R}$ , with the signature of  $\ell$ -monoids with designated constants  $-1$  and  $1$ , i.e.  $\{+, \vee, \wedge, 0, 1, -1\}$ .

Hypotheses on the generating algebra  $\mathbf{L}$  (think e.g. of  $\mathbb{R}$ ):

1.  $\mathbf{L}$  has a **majority term**. (E.g.:  $\mathbf{L}$  has a lattice reduct.) (It can be generalized to near-unanimity terms.)
2. ( $\mathbf{L}$  is “**indecomposable**”:)  $\mathbf{L}$  is *hereditarily finitely subdirectly irreducible*, i.e. every subalgebra of  $\mathbf{L}$  is finitely subdirectly irreducible. (E.g.: every subalgebra of  $\mathbf{L}$  is simple, and  $\mathbf{L}$  has two distinct constants.)
3. ( $\mathbf{L}$  is “**rigid**”:) For each subalgebra  $\mathbf{A}$  of  $\mathbf{L}$ , the inclusion  $\mathbf{A} \hookrightarrow \mathbf{L}$  is the unique homomorphism from  $\mathbf{A}$  to  $\mathbf{L}$ .



Under these hypotheses, we provide a duality for the category of algebras in  $\mathbf{ISP}(\mathbf{L})$  that are *finitely  $\mathbf{L}$ -valued*.

We will represent an algebra via a

1. Stone space  $X$ , together with
2. a “local constraint” for each subset  $I \subseteq X$  of cardinality  $\leq 2$ .

For Priestley duality, {constraints} = order.


## Example

$\mathbf{L} = \langle \mathbb{R}; +, \vee, \wedge, 0, 1, -1 \rangle$ . (Commutative lattice-ordered monoid...)

$$\mathbf{A} := \{f: \mathbb{N} \rightarrow \mathbb{R} \mid f \text{ is eventually constant}\}.$$

$\mathbf{A} \stackrel{?}{\cong} \{\text{cont. functions over a Stone space satisfying local constraints}\}.$


Stone space:  $\alpha\mathbb{N} = \mathbb{N} \cup \{\infty\}$ , the one-point compactification of  $\mathbb{N}$ .



A continuous function from  $\alpha\mathbb{N}$  to the discrete space  $\mathbb{R}$ .

$\mathbf{A} \cong$  algebra of continuous functions from  $\alpha\mathbb{N}$  to the discrete space  $\mathbb{R}$ .  
(No local constraints.)

$\mathbf{B} := \{f: \mathbb{N} \rightarrow \mathbb{R} \mid f \text{ is eventually constant and order-preserving}\}.$



A continuous order-preserving function from  $\alpha\mathbb{N}$  to the discrete space  $\mathbb{R}$ .

$\mathbf{B} \cong$  algebra of *order-preserving* continuous functions from  $\alpha\mathbb{N}$  to the discrete space  $\mathbb{R}$ .

*Order-preservation* is given by a family of *binary constraints*.

Constraint on  $\{2, 3\}$ :  $\{(x, y) \in \mathbb{R}^2 \mid x \leq y\}.$

We equip  $\mathbf{L}$  with the discrete topology. A continuous function from a Stone space  $X$  to  $\mathbf{L}$  has finite image.

" $I \subseteq_2 X$ " stands for " $I$  is a subset of  $X$  of cardinality at most 2".

## Definition

A Priestley  $\mathbf{L}$ -space consists of a Stone space  $X$  and, for each  $I \subseteq_2 X$ , of a subalgebra  $\mathbf{A}_I$  of  $\mathbf{L}^I$  s.t.

1. (**Local-to-global extension**) For all  $I \subseteq_2 X$ , every  $f \in \mathbf{A}_I$  has a continuous extension  $g: X \rightarrow \mathbf{L}$  that satisfies all constraints, i.e. s.t., for all  $J \subseteq_2 X$ ,  $g|_J \in \mathbf{A}_J$ .
2. (**Separation**) For all distinct  $x, y \in X$ , there is  $f \in \mathbf{A}_{\{x,y\}}$  s.t.  $f(x) \neq f(y)$ .

E.g.:  $X = \alpha\mathbb{N}$ . For  $I \subseteq_2 \alpha\mathbb{N}$ ,  $\mathbf{A}_I := \{f: I \rightarrow \mathbb{R} \mid f \text{ is order-preserving}\}$ .

## Definition

An algebra  $\mathbf{A}$  in  $\mathbf{ISP}(\mathbf{L})$  is said to be *finitely  $\mathbf{L}$ -valued* if for each  $a \in \mathbf{A}$  the set

$$\{h(a) : h : \mathbf{A} \rightarrow \mathbf{L} \text{ homomorphism}\}$$

is finite.

I.e., each element of  $\mathbf{A}$ , thought of as a function  $X \rightarrow \mathbf{L}$ , has finite image.

Example: For  $\mathbf{L} = \langle \mathbb{R}; +, \vee, \wedge, 0, 1, -1 \rangle$ , the following are finitely  $\mathbf{L}$ -valued:

- ▶  $\mathbb{R}$ ,
- ▶ any finite power  $\mathbb{R}^n$  of  $\mathbb{R}$ ,
- ▶ any subalgebra of a finite power  $\mathbb{R}^n$  of  $\mathbb{R}$ ,
- ▶  $\{f : \mathbb{N} \rightarrow \mathbb{R} \mid f \text{ is eventually constant}\}$ .

Hypotheses on  $\mathbf{L}$  (think of  $2$ , or  $[0, 1]$ , or  $\mathbb{R}$ ):

1.  $\mathbf{L}$  has a majority term. (E.g.:  $\mathbf{L}$  has a lattice reduct.)
2. ( $\mathbf{L}$  is “indecomposable”:)  $\mathbf{L}$  is *hereditarily finitely subdirectly irreducible*, i.e. every subalgebra of  $\mathbf{L}$  is finitely subdirectly irreducible. (E.g.: every subalgebra of  $\mathbf{L}$  is simple, and  $\mathbf{L}$  has two distinct constants.)
3. ( $\mathbf{L}$  is “rigid”:) For each subalgebra  $\mathbf{A}$  of  $\mathbf{L}$ , the inclusion  $\mathbf{A} \hookrightarrow \mathbf{L}$  is the unique homomorphism from  $\mathbf{A}$  to  $\mathbf{L}$ .

## Theorem (Main result)

Suppose  $\mathbf{L}$  satisfies (1–3). The category of finitely  $\mathbf{L}$ -valued algebras in  $\text{ISP}(\mathbf{L})$  (and homomorphisms) is dually equivalent to the category of Priestley  $\mathbf{L}$ -spaces (and appropriate morphisms).

Thank you!